

Adamax

PEPTIDES

SPECIAL HANDLING/STORAGE SPECIFICATIONS None ADDITIONAL ITEMS REQUIRED

Bacteriostatic water Recommended insulin syringes for dosing: 1mL-30 gauge-8mm (5/16") or .5mL-30 gauge-8mm (5/16") GENERAL DOSING GUIDELINES 8-12 week cycle followed by washout period equal to the length of t

SPECIAL HANDLING/STORAGE SPECIFICATIONS None ADDITIONAL ITEMS REQUIRED

Bacteriostatic water Recommended insulin syringes for dosing: 1mL-30 gauge-8mm (5/16") or .5mL-30 gauge-8mm (5/16") GENERAL DOSING GUIDELINES 8-12 week cycle followed by washout period equal to the length of the cycle you completed Daily Dosing Morning dosing recommended, although not absolutely mandatory; try to be consistent with your dosing time each day You do not have to be fasted to dose Subcutaneous Administration RECONSTITUTION Mix with 3mL (300 units) of bacteriostatic water DOSING PROTOCOL Weeks 1-2: Take 9 units (300mcg) once per day Weeks 3-4: Take 15 units (500mcg) once per day Weeks 5-6: Take 23 units (750mcg) once per day Weeks 7-8+: Take 30 units (1,000mcg) once per day

RESEARCH USE DISCLAIMER: All information in this document is provided strictly for laboratory research and educational purposes. It is not intended for human or veterinary use, diagnosis, or treatment.