

Ara-290

PEPTIDES

ARA-290 (aka Cibinetide) is a synthetic peptide derived from erythropoietin (EPO) designed to activate the innate repair receptor (IRR) and produce anti-inflammatory, neuroprotective, and tissue-repair effects without EPO's red blood-cell-stimulating or pro-thrombotic actions. Ben

ARA-290 (aka Cibinetide) is a synthetic peptide derived from erythropoietin (EPO) designed to activate the innate repair receptor (IRR) and produce anti-inflammatory, neuroprotective, and tissue-repair effects without EPO's red blood-cell-stimulating or pro-thrombotic actions. Benefits Relief of neuropathic pain (especially small-fiber neuropathy) Strong anti-inflammatory Supports nerve repair/regeneration Restores immune balance in autoimmune disorders Tissue repair actions in injured, stressed, or ischemic tissues Provides adjunctive support for metabolic diseases like Type 2 diabetes, helping nerve complications, lipid/glucose improvements Does not produce clinically significant changes in electrolytes, liver, kidney, pancreas, erythrocytes, platelets, or leukocytes Improves lipid and glucose metabolism in diabetics Interactions Concomitant use of strong immunosuppressants or biologic anti-inflammatory drugs may amplify immune modulation (caution advised) Agents that increase angiogenesis or that affect repair/vascular pathways may theoretically have additive or synergistic effects Given the repair/angiogenic axis, close monitoring of anticoagulant and antiplatelet medications is recommended Concomitant use of antidiabetic agents may require dose monitoring due to a risk of hypoglycemia Contraindications Do not use if you are pregnant or breastfeeding Do not use if you determine that you are hypersensitive to this peptide Do not use if you have an active malignancy Do not use if you have uncontrolled cardiovascular disease or are at high risk of thrombosis ARA-290: 16MG RECONSTITUTION AND DOSING 4, 8, or 12 week cycles followed by 2, 4, or 6 week washout periods Subcutaneous Administration Standard Protocol Mix with 1.5mL (150 units) of BAC water 18, 28, or 38 units (2, 3, or 4mg), 1x/day, daily Maximum Protocol Mix with 1mL (150 units) of BAC water 75 units (8mg), 1x/day, daily, for a maximum of 4 weeks

RESEARCH USE DISCLAIMER: All information in this document is provided strictly for laboratory research and educational purposes. It is not intended for human or veterinary use, diagnosis, or treatment.