

BPC-157-TB-500-Cartalax

PEPTIDES

BPC-157: 5mg TB-500: 5mg Cartalax: 10mg CYCLE LENGTH AND WASHOUT PERIOD: 8 week cycle followed by 4 week washout period (Must have a cycle and washout period because of the inclusion of the TB-500 and Cartalax) DOSING TIME: Any time of day or night FASTED DOSING?: You do not have

BPC-157: 5mg TB-500: 5mg Cartalax: 10mg CYCLE LENGTH AND WASHOUT PERIOD: 8 week cycle followed by 4 week washout period (Must have a cycle and washout period because of the inclusion of the TB-500 and Cartalax) DOSING TIME: Any time of day or night FASTED DOSING?: You do not have to be in a fasted state to dose. ROUTE OF ADMINISTRATION: Subcutaneous injection. RECONSTITUTION SOLUTION: Bacteriostatic water DOSING Mix with 2mL (200 units) of Bacteriostatic water STANDARD PROTOCOL Weeks 1-2: 6 units, 1x/day Weeks 3-8: 8 units, 1x/day MAXIMUM PROTOCOL Weeks 1-2: 10 units, 1x/day Weeks 3-8: 6 units, 1x/day

RESEARCH USE DISCLAIMER: All information in this document is provided strictly for laboratory research and educational purposes. It is not intended for human or veterinary use, diagnosis, or treatment.