

Hulk-2

PEPTIDES

“The Hulk:” Blend DOSING: CYCLE, ADMINISTRATION, AND GENERAL CONSIDERATIONS 12-16 week cycle followed by 8-16 week washout period Dose once per day, daily, for the duration of the cycle Best time to dose is immediately before bedtime, fasted for 2+ hours; may inject in the mornin

“The Hulk:” Blend DOSING: CYCLE, ADMINISTRATION, AND GENERAL CONSIDERATIONS 12-16 week cycle followed by 8-16 week washout period Dose once per day, daily, for the duration of the cycle Best time to dose is immediately before bedtime, fasted for 2+ hours; may inject in the morning upon wake-up prior to food if preferred Subcutaneous injection Inject slowly to avoid hot, flushing, pins and needles sensation during and following injection; after you draw your dose, you may consider drawing up 20 or 30 units of BAC water to dilute it with, making it easier to slowly introduce the peptide to your system. 30 or 50 unit syringes (.3 or .5mL) syringes are highly recommended: each unit on these syringes has its own line, allowing the accurate measurement of the small doses that this blend calls for RECONSTITUTION Mix with 2.5mL (250 units) of BAC water DOSING Weeks 1-2: 1 units (100mcg), 1x/day, daily Weeks 3-4: 2 units (150mcg), 1x/day, daily Weeks 5-6: 3 units (200mcg), 1x/day, daily Weeks 7-12: 4 units (250mcg), 1x/day, daily Weeks 12-16 Optional: 5 units (300mcg), 1x/day, daily : Yes, we are aware that many of the “actual” unit calculations work out to a partial unit. Because a single unit is often difficult to accurately measure on a syringe, the idea of directing someone to attempt to measure out a half unit was ludicrous, and I took the liberty of rounding the answer. This is a low dose protocol for anti-aging. For bodybuilders and/or those who wish to achieve maximum results, increase your dose

RESEARCH USE DISCLAIMER: All information in this document is provided strictly for laboratory research and educational purposes. It is not intended for human or veterinary use, diagnosis, or treatment.